

Two kinds of conditionings for stable Lévy processes

Kouji YANO

Abstract

Two kinds of conditionings for one-dimensional stable Lévy processes are discussed via h -transforms of excursion measures: One is to stay positive, and the other is to avoid the origin.

1 Introduction

It is well-known that a one-dimensional Brownian motion conditioned to stay positive is a three-dimensional Bessel process. As an easy consequence, it follows that the former conditioned to avoid the origin is a symmetrized one of the latter.

The aim of the present article is to give a brief survey on these two different kinds of conditionings for one-dimensional stable Lévy processes via h -transforms of Itô's excursion measures.

The organization of this article is as follows. In Section 2, we recall the conditionings for Brownian motions. In Section 3, we give a review on the conditioning for stable Lévy processes to stay positive. In Section 4, we present results on the conditioning for symmetric stable Lévy processes to avoid the origin.

2 Conditionings for Brownian motions

We recall the conditionings for Brownian motions. For the details, see, e.g., [8, §III.4.3] and [10, §VI.3 and Chap.XII].

Let (X_t) denote the coordinate process on the space of càdlàg functions and $(\mathcal{F}_t)$ its natural filtration. Set $\mathcal{F}_\infty = \sigma(\cup_{t>0} \mathcal{F}_t)$. For $t \geq 0$, we write θ_t for the shift operator: $X_s \circ \theta_t = X_{t+s}$. For $0 < t < \infty$, a functional Z_t is called $\mathcal{F}_t$ -nice if Z_t is of the form $Z_t = f(X_{t_1}, \dots, X_{t_n})$ for some $0 < t_1 < \dots < t_n < t$ and some continuous function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ which vanishes at infinity.

Let W_x denote the law of the one-dimensional Brownian motion starting from $x \in \mathbb{R}$.

2.1 Brownian motions conditioned to stay positive

For any fixed $t > 0$, we define a probability law $W^{\uparrow, (t)}$ on $\mathcal{F}_t$ as

$$W^{\uparrow, (t)}(\cdot) = \frac{n^+(\cdot; \zeta > t)}{n^+(\zeta > t)} \quad (2.1)$$

Department of Mathematics, Graduate School of Science, Kobe University, Kobe, Japan.
Research supported by KAKENHI (20740060)

where $\mathbf{n}^+$ stands for the *excursion measure of the reflecting Brownian motion* (see, e.g., [8, §III.4.3] and [10, §§XII.4]) and ζ for the *lifetime*. The process $(X_s : s \leq t)$ under $W^{\uparrow, (t)}$ is called the *Brownian meander*. Durrett–Iglehart–Miller [7, Thm.2.1] have proved that

$$W^{\uparrow, (t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} W_0 \left[Z_t \mid \forall u \leq t, X_u \geq -\varepsilon \right] \quad (2.2)$$

for any bounded continuous $\mathcal{F}_t$ -measurable functional Z_t ; in particular, for any $\mathcal{F}_t$ -nice functional Z_t . We may represent (2.2) symbolically as

$$W^{\uparrow, (t)}(\cdot) = W_0 \left(\cdot \mid \forall u \leq t, X_u \geq 0 \right); \quad (2.3)$$

that is, the Brownian meander is the Brownian motion *conditioned to stay positive until time t* . As another interpretation of (2.3), we have

$$W^{\uparrow, (t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} W_0 \left[Z_t \circ \theta_\varepsilon \mid \forall u \leq t, X_u \circ \theta_\varepsilon \geq 0 \right] \quad (2.4)$$

for any $\mathcal{F}_t$ -nice functional Z_t . The proof of (2.4) will be given in Theorem 3.5 in the settings of stable Lévy processes.

We write $W_0^\uparrow$ for the law of a *three-dimensional Bessel process*, that is, the law of the radius $\sqrt{(B_t^{(1)})^2 + (B_t^{(2)})^2 + (B_t^{(3)})^2}$ of a three-dimensional Brownian motion $(B_t^{(1)}, B_t^{(2)}, B_t^{(3)})$. We remark that $W_0^\uparrow$ is locally equivalent to $\mathbf{n}^+$:

$$dW_0^\uparrow|_{\mathcal{F}_t} = \frac{1}{C^\uparrow} X_t d\mathbf{n}^+|_{\mathcal{F}_t} \quad (2.5)$$

where $C^\uparrow = \mathbf{n}^+[X_t]$ is a constant independent of $t > 0$. We say that $W_0^\uparrow$ is the *h -transform of the excursion measure $\mathbf{n}^+$ with respect to the function $h(x) = x$* . Then it holds (see, e.g., Theorem 3.6) that

$$W_0^\uparrow[Z] = \lim_{t \rightarrow \infty} W^{\uparrow, (t)}[Z] \quad (2.6)$$

for any $\mathcal{F}_t$ -nice functional Z with $0 < t < \infty$. We may represent (2.6) symbolically as

$$W_0^\uparrow(\cdot) = W_0 \left(\cdot \mid \forall u, X_u \geq 0 \right); \quad (2.7)$$

that is, $W_0^\uparrow$ is the Brownian motion *conditioned to stay positive during the whole time*.

2.2 Brownian motions conditioned to avoid the origin

For any fixed $t > 0$, we define a probability law $W^{\times, (t)}$ on $\mathcal{F}_t$ as

$$W^{\times, (t)} = \frac{W^{\uparrow, (t)} + W^{\downarrow, (t)}}{2} \quad (2.8)$$

where $W^{\downarrow,(t)}$ stands for the law of the process $(-X_s : s \leq t)$ under $W^{\uparrow,(t)}$. The law $W^{\times,(t)}$ may be represented as

$$W^{\times,(t)}(\cdot) = \frac{\mathbf{n}(\cdot; \zeta > t)}{\mathbf{n}(\zeta > t)} \quad (2.9)$$

where $\mathbf{n}$ stands for the *excursion measure of the Brownian motion*; in fact, $\mathbf{n} = \frac{\mathbf{n}^+ + \mathbf{n}^-}{2}$ where $\mathbf{n}^-$ is the image measure of the process $(-X_t)$ under $\mathbf{n}^+$. Immediately from (2.4) and continuity of paths, we have

$$W^{\times,(t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} W_0 \left[Z_t \circ \theta_\varepsilon \mid \forall u \leq t, X_u \circ \theta_\varepsilon \neq 0 \right] \quad (2.10)$$

for any $\mathcal{F}_t$ -nice functional Z_t . We may represent (2.10) symbolically as

$$W^{\times,(t)}(\cdot) = W_0 \left(\cdot \mid \forall u \leq t, X_u \neq 0 \right); \quad (2.11)$$

that is, $W^{\times,(t)}$ is the Brownian motion *conditioned to avoid the origin until time t* .

We define

$$W_0^\times = \frac{W_0^\uparrow + W_0^\downarrow}{2} \quad (2.12)$$

where $W_0^\downarrow$ stands for the law of the process $(-X_t)$ under $W_0^\uparrow$. In other words, the law $W_0^\uparrow$ is the symmetrization of the three-dimensional Bessel process. We also remark that $W_0^\times$ is locally equivalent to $\mathbf{n}$:

$$dW_0^\times|_{\mathcal{F}_t} = \frac{1}{C^\times} |X_t| d\mathbf{n}|_{\mathcal{F}_t} \quad (2.13)$$

where $C^\times = \mathbf{n}[|X_t|]$ is a constant independent of $t > 0$. We say that $W_0^\times$ is the *h -transform with respect to the function $h(x) = |x|$* . Then it is immediate from (2.6) that

$$W_0^\times[Z] = \lim_{t \rightarrow \infty} W^{\times,(t)}[Z] \quad (2.14)$$

for any $\mathcal{F}_t$ -nice functional Z with $0 < t < \infty$. We may represent (2.14) symbolically as

$$W_0^\times(\cdot) = W_0 \left(\cdot \mid \forall u, X_u \neq 0 \right); \quad (2.15)$$

that is, $W_0^\times$ is the Brownian motion *conditioned to avoid the origin during the whole time*.

3 Stable Lévy processes conditioned to stay positive

Let us review the theory of strictly stable Lévy processes conditioned to stay positive. For concise references, see, e.g., [1] and [6]. We refer to these textbooks also about the theory of conditioning to stay positive for *spectrally negative Lévy processes*, where we do not go into the details.

For a Borel set F , we denote the first hitting time of F by

$$T_F = \inf\{t > 0 : X_t \in F\}. \quad (3.1)$$

Define

$$\underline{X}_t = \inf_{s \leq t} X_s, \quad \underline{R}_t = X_t - \underline{X}_t \quad (3.2)$$

and call the process $(\underline{R}_t)$ the *reflected process*.

Let (P_x) denote the law of a *strictly stable Lévy process of index* $0 < \alpha \leq 2$, that is, a process with càdlàg paths and with stationary independent increments satisfying the following scaling property:

$$(k^{-\frac{1}{\alpha}} X_{kt} : t \geq 0) \stackrel{\text{law}}{=} (X_t : t \geq 0) \quad \text{under } P_0 \quad (3.3)$$

for any $k > 0$. Note that the Brownian case corresponds to $\alpha = 2$. From the scaling property (3.3), it is immediate that the quantity

$$\rho := P_0(X_t \geq 0) \quad (3.4)$$

does not depend on $t > 0$, which is called the *positivity parameter*. The possible values of ρ range over $[0, 1]$ if $0 < \alpha < 1$, $(0, 1)$ if $\alpha = 1$, and $[1 - \frac{1}{\alpha}, \frac{1}{\alpha}]$ if $1 < \alpha \leq 2$. Let $(Q_x : x > 0)$ denote the law of the process killed at $T_{(-\infty, 0)}$:

$$Q_x(\Lambda_t; t < \zeta) = P_x(\Lambda_t; t < T_{(-\infty, 0)}), \quad x > 0, \Lambda_t \in \mathcal{F}_t. \quad (3.5)$$

Note that the function

$$(x, t) \mapsto Q_x(t < \zeta) = P_x(t < T_{(-\infty, 0)}) \quad (3.6)$$

is jointly continuous in $x > 0$ and $t > 0$.

Let us **exclude** the case where $|X|$ is a subordinator, i.e.,

$$0 < \alpha < 1 \quad \text{and} \quad \rho = 0, 1. \quad (3.7)$$

Then the reflected process $(\underline{R}_t)$ under (P_x) is a Feller process where the origin is regular for itself, and hence there exists the continuous local time process $(\underline{L}_t)$ at level 0 of the reflected process $(\underline{R}_t)$ such that

$$P_0 \left[\int_0^\infty e^{-qt} d\underline{L}_t \right] = q^{\rho-1}, \quad q > 0. \quad (3.8)$$

Let $\mathbf{n}^\uparrow$ denote the corresponding excursion measure away from 0 of the reflected process $(\underline{R}_t)$. The Markov property of $\mathbf{n}^\uparrow$ may be expressed as

$$\mathbf{n}^\uparrow(1_\Lambda \circ \theta_t; \Lambda_t, t < \zeta) = \mathbf{n}^\uparrow[Q_{X_t}(\Lambda); \Lambda_t, t < \zeta] \quad (3.9)$$

for any $\Lambda \in \mathcal{F}_\infty$ and $\Lambda_t \in \mathcal{F}_t$. We introduce the following function:

$$h^\uparrow(x) = P_0 \left[\int_0^\infty 1_{\{\underline{X}_t \geq -x\}} d\underline{L}_t \right], \quad x \geq 0. \quad (3.10)$$

Since the *ladder height process* $H := \underline{X} \circ \underline{L}^{-1}$ is a stable Lévy process of index $\alpha(1 - \rho)$, we see that

$$h^\uparrow(x) = P_0 \left[\int_0^\infty 1_{\{H_u \geq -x\}} du \right] = C_1^\uparrow x^{\alpha(1-\rho)} \quad (3.11)$$

for some constant $C_1^\uparrow > 0$ independent of $x \geq 0$. The following theorem is due to Silverstein [14, Thm.2]; another proof can be found in Chaumont–Doney [5, Lem.1] (see also Doney [6, Lem.10 in §8.3]).

Theorem 3.1 ([14]). *It holds that*

$$Q_x[(X_t)^{\alpha(1-\rho)}; t < \zeta] = x^{\alpha(1-\rho)}, \quad x > 0, t > 0, \quad (3.12)$$

$$\mathbf{n}^\uparrow[(X_t)^{\alpha(1-\rho)}; t < \zeta] = C_2^\uparrow, \quad t > 0 \quad (3.13)$$

for some constant $C_2^\uparrow$ independent of $t > 0$.

By virtue of this theorem, we may define the h -transform by

$$dP_x^\uparrow|_{\mathcal{F}_t} = \begin{cases} \left(\frac{X_t}{x}\right)^{\alpha(1-\rho)} dQ_x|_{\mathcal{F}_t} & \text{if } x > 0, \\ \frac{1}{C_2^\uparrow} (X_t)^{\alpha(1-\rho)} d\mathbf{n}^\uparrow|_{\mathcal{F}_t} & \text{if } x = 0; \end{cases} \quad (3.14)$$

indeed, the family $(P_x^\uparrow|_{\mathcal{F}_t} : t \geq 0)$ is proved to be consistent by the Markov property of $\mathbf{n}^\uparrow$.

Theorem 3.2 ([3]). *The process $((X_t), (P_x^\uparrow))$ is a Feller process.*

This theorem is due to Chaumont [3, Thm.6], where he proved weak convergence $P_x^\uparrow \rightarrow P_0^\uparrow$ as $x \rightarrow 0+$ in the càdlàg space equipped with Skorokhod topology. Bertoin–Yor [2, Thm.1] proved the weak convergence for general positive self-similar Markov processes. Tanaka [15, Thm.4] proved the Feller property for quite a general class of Lévy processes.

Now let us discuss the conditionings (2.3) and (2.7) for stable Lévy processes. The following theorem is an immediate consequence of Chaumont [4, Lem.1] and of the continuity of (3.6).

Theorem 3.3 ([4]). *Let $t > 0$ be fixed. Then the function*

$$[0, \infty) \ni x \mapsto P_x^\uparrow [(X_t)^{-\alpha(1-\rho)}] \quad (3.15)$$

is continuous and vanishes at infinity.

Define a probability law $M^{\uparrow, (t)}$ on $\mathcal{F}_t$ as

$$M^{\uparrow, (t)}(\Lambda_t) = \frac{\mathbf{n}^\uparrow(\Lambda_t; \zeta > t)}{\mathbf{n}^\uparrow(\zeta > t)} = \frac{P_0^\uparrow[(X_t)^{-\alpha(1-\rho)}; \Lambda_t]}{P_0^\uparrow[(X_t)^{-\alpha(1-\rho)}]} \quad (3.16)$$

for $\Lambda_t \in \mathcal{F}_t$. The following theorem, which generalizes (2.2), can be found in Bertoin [1, Thm.VIII.18].

Theorem 3.4 ([1]). *For any $t > 0$, it holds that*

$$M^{\uparrow, (t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} P_0 \left[Z_t \mid \forall u \leq t, X_u \geq -\varepsilon \right] \quad (3.17)$$

for any $\mathcal{F}_t$ -nice functional Z_t .

Now we give the following version of (2.4) for stable Lévy processes.

Theorem 3.5. *For any $t > 0$, it holds that*

$$M^{\uparrow, (t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} P_0 \left[Z_t \circ \theta_\varepsilon \mid \forall u \leq t, X_u \circ \theta_\varepsilon \geq 0 \right] \quad (3.18)$$

for any $\mathcal{F}_t$ -nice functional Z_t .

Proof. By the Markov property, the expectation of the right hand side of (3.18) is equal to

$$\frac{P_0 [P_{X_\varepsilon} [Z_t; \forall u \leq t, X_u \geq 0]]}{P_0 [P_{X_\varepsilon} (\forall u \leq t, X_u \geq 0)]} = \frac{P_0 [(X_\varepsilon)^{-\gamma} P_{X_\varepsilon}^\uparrow [Z_t(X_t)^{-\gamma}]]}{P_0 [(X_\varepsilon)^{-\gamma} P_{X_\varepsilon}^\uparrow [(X_t)^{-\gamma}]]}, \quad (3.19)$$

where we put $\gamma = \alpha(1 - \rho)$. By the scaling property, this is equal to

$$\frac{P_0 [(X_1)^{-\gamma} P_{\varepsilon^{1/\alpha} X_1}^\uparrow [Z_t(X_t)^{-\gamma}]]}{P_0 [(X_1)^{-\gamma} P_{\varepsilon^{1/\alpha} X_1}^\uparrow [(X_t)^{-\gamma}]]}. \quad (3.20)$$

By Theorems 3.2 and 3.3 and by the dominated convergence theorem, we see that this quantity converges as $\varepsilon \rightarrow 0+$ to

$$\frac{P_0^\uparrow [Z_t(X_t)^{-\gamma}]}{P_0^\uparrow [(X_t)^{-\gamma}]}, \quad (3.21)$$

which coincides with $M^{\uparrow, (t)}[Z_t]$ by the definition (3.16). □

The following theorem generalizes (2.6).

Theorem 3.6. *It holds that*

$$P_0^\uparrow[Z] = \lim_{t \rightarrow \infty} M^{\uparrow, (t)}[Z] \quad (3.22)$$

for any $\mathcal{F}_t$ -nice functional Z with $0 < t < \infty$.

The proof can be done in the same way as Theorem 3.5.

4 Symmetric stable Lévy processes conditioned to avoid the origin

Let us discuss the conditioning for symmetric stable Lévy processes to avoid the origin. This has been introduced by Yano–Yano–Yor [18] in order to extend some of the *penalisation problems* for Brownian motions by Roynette–Vallois–Yor [11], [12] and Najnudel–Roynette–Yor [9], to symmetric stable Lévy processes.

We assume that $\rho = \frac{1}{2}$, i.e., the process $((X_t), (P_x))$ is *symmetric*, and that the index α satisfies $1 < \alpha \leq 2$. For simplicity, we assume that $P_0[e^{i\lambda X_1}] = e^{-|\lambda|^\alpha}$. Then the origin is regular for itself, and there exists the continuous resolvent density:

$$u_q(x) = \frac{1}{\pi} \int_0^\infty \frac{\cos x\lambda}{q + \lambda^\alpha} d\lambda, \quad q > 0, \quad x \in \mathbb{R}. \quad (4.1)$$

Moreover, there exists the local time process at level 0 of the process (X_t) , which we denote by (L_t) , such that

$$P_0 \left[\int_0^\infty e^{-qt} dL_t \right] = u_q(0) = \frac{q^{\frac{1}{\alpha}-1}}{\alpha \sin \frac{\pi}{\alpha}}, \quad q > 0. \quad (4.2)$$

The corresponding excursion measure away from 0 will be denoted by $\mathbf{n}^\times$. We introduce the following function:

$$h^\times(x) = \lim_{q \rightarrow 0+} \{u_q(0) - u_q(x)\} = \frac{1}{C^\times} |x|^{\alpha-1}, \quad x \in \mathbb{R}; \quad (4.3)$$

see, e.g., [17, Appendix] for the exact value of the constant $C^\times$. Note that the function $h^\times(x)$ may also be represented as

$$h^\times(x) = P_0[L_{T_{\{x\}}}], \quad x \in \mathbb{R}; \quad (4.4)$$

see, e.g., [1, Lem.V.11]. Let (P_x^0) denote the law of the process $((X_t), (P_x))$ killed at $T_{\{0\}}$:

$$P_x^0(\Lambda_t; t < \zeta) = P_x(\Lambda_t; t < T_{\{0\}}), \quad x \neq 0, \quad \Lambda_t \in \mathcal{F}_t. \quad (4.5)$$

By [18, Thm.3.5], we see that the function

$$(x, t) \mapsto P_x^0(t < \zeta) = P_x(t < T_{\{0\}}) \quad (4.6)$$

is jointly continuous in $x \in \mathbb{R} \setminus \{0\}$ and $t > 0$. The following theorem is due to Salminen–Yor [13, eq.(3)] and Yano–Yano–Yor [18, Thm.4.7].

Theorem 4.1 ([13], [18]). *It holds that*

$$P_x^0[|X_t|^{\alpha-1}] = |x|^{\alpha-1}, \quad x \neq 0, \quad t > 0, \quad (4.7)$$

$$\mathbf{n}^\times[|X_t|^{\alpha-1}; t < \zeta] = C^\times, \quad t > 0. \quad (4.8)$$

By virtue of this theorem, we may define the h -transform by

$$dP_x^\times|_{\mathcal{F}_t} = \begin{cases} \left|\frac{X_t}{x}\right|^{\alpha-1} dP_x^0|_{\mathcal{F}_t} & \text{if } x \neq 0, \\ \frac{1}{C^\times} |X_t|^{\alpha-1} d\mathbf{n}^\times|_{\mathcal{F}_t} & \text{if } x = 0; \end{cases} \quad (4.9)$$

indeed, the family $(P_x^\times|_{\mathcal{F}_t} : t \geq 0)$ is proved to be consistent by the Markov property of $\mathbf{n}^\times$. The following theorem is due to [16, Thm.1.5].

Theorem 4.2 ([16]). *Suppose that $1 < \alpha < 2$. Then the process $((X_t), (P_x^\times))$ is a Feller process.*

Remark 4.3. Yano [16, Thm.1.4 and Cor.1.9] obtained the following long-time behavior of paths: If $1 < \alpha < 2$, then

$$P_0^\times \left(\limsup_{t \rightarrow \infty} X_t = \limsup_{t \rightarrow \infty} (-X_t) = \lim_{t \rightarrow \infty} |X_t| = \infty \right) = 1. \quad (4.10)$$

Remark 4.4. In the Brownian case ($\alpha = 2$), the process $((X_t), (P_x^\times))$ is *not* a Feller process. Indeed, $P_0^\times$ is not irreducible (see (2.12)). Contrary to (4.10), the long-time behavior in this case is as follows:

$$P_0^\times \left(\lim_{t \rightarrow \infty} X_t = \infty \right) = P_0^\times \left(\lim_{t \rightarrow \infty} X_t = -\infty \right) = \frac{1}{2}. \quad (4.11)$$

Now let us discuss the conditionings (2.11) and (2.15) for symmetric stable Lévy processes. The following theorem is an immediate consequence of Yano–Yano–Yor [18, Lem.4.10] and of the continuity of (4.6).

Theorem 4.5 ([18]). *Let $t > 0$ be fixed. Then the function*

$$\mathbb{R} \ni x \mapsto P_x^\times [|X_t|^{-(\alpha-1)}] \quad (4.12)$$

is continuous and vanishes at infinity.

Define a probability law $M^{\times,(t)}$ on $\mathcal{F}_t$ as

$$M^{\times,(t)}(\Lambda_t) = \frac{\mathbf{n}^\times(\Lambda_t; \zeta > t)}{\mathbf{n}^\times(\zeta > t)} = \frac{P_0^\times[|X_t|^{-(\alpha-1)}; \Lambda_t]}{P_0^\uparrow[|X_t|^{-(\alpha-1)}]} \quad (4.13)$$

for $\Lambda_t \in \mathcal{F}_t$. The following theorem generalizes (2.10).

Theorem 4.6. *For any $t > 0$, it holds that*

$$M^{\times,(t)}[Z_t] = \lim_{\varepsilon \rightarrow 0+} P_0 \left[Z_t \circ \theta_\varepsilon \mid \forall u \leq t, X_u \circ \theta_\varepsilon \neq 0 \right] \quad (4.14)$$

for any $\mathcal{F}_t$ -nice functional Z_t .

The proof can be done in the same way as Theorem 3.5 by virtue of Theorems 4.2 and 4.5. The following theorem generalizes (2.14).

Theorem 4.7 ([18]). *It holds that*

$$P_0^\times[Z] = \lim_{t \rightarrow \infty} M^{\times,(t)}[Z] \quad (4.15)$$

for any $\mathcal{F}_t$ -nice functional Z with $0 < t < \infty$.

This is a special case of [18, Thm.4.9].

Acknowledgements. The author thanks Professor Marc Yor for valuable comments.

References

- [1] J. Bertoin. *Lévy processes*. Cambridge University Press, Cambridge, 1996.
- [2] J. Bertoin and M. Yor. The entrance laws of self-similar Markov processes and exponential functionals of Lévy processes. *Potential Anal.*, 17(4):389–400, 2002.
- [3] L. Chaumont. Conditionings and path decompositions for Lévy processes. *Stochastic Process. Appl.*, 64(1):39–54, 1996.
- [4] L. Chaumont. Excursion normalisée, méandre et pont pour les processus de Lévy stables. *Bull. Sci. Math.*, 121(5):377–403, 1997.
- [5] L. Chaumont and R. A. Doney. On Lévy processes conditioned to stay positive. *Electron. J. Probab.*, 10:no. 28, 948–961 (electronic), 2005; corrections in 13:no. 1, 1–4 (electronic), 2008.
- [6] R. A. Doney. *Fluctuation theory for Lévy processes*. LNM 1897. Springer, Berlin, 2007. Lectures from the 35th Summer School in Saint-Flour, 2005.
- [7] R. T. Durrett, D. L. Iglehart, and D. R. Miller. Weak convergence to Brownian meander and Brownian excursion. *Ann. Probab.*, 5(1):117–129, 1977.
- [8] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*. North-Holland Publishing Co., Amsterdam, 2nd edition, 1989.
- [9] J. Najnudel, B. Roynette, and M. Yor. A remarkable σ -finite measure on $\mathcal{C}(\mathbf{R}_+, \mathbf{R})$ related to many Brownian penalisations. *C. R. Math. Acad. Sci. Paris*, 345(8):459–466, 2007.
- [10] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*. Springer-Verlag, Berlin, 3rd edition, 1999.
- [11] B. Roynette, P. Vallois, and M. Yor. Some penalisations of the Wiener measure. *Jpn. J. Math.*, 1(1):263–290, 2006.
- [12] B. Roynette and M. Yor. Penalising Brownian paths. LNM 1969. Springer, forthcoming, 2009.
- [13] P. Salminen and M. Yor. Tanaka formula for symmetric Lévy processes. In *Sém. de Probab. XL*, LNM 1899: 265–285. Springer, Berlin, 2007.
- [14] M. L. Silverstein. Classification of coharmonic and coinvariant functions for a Lévy process. *Ann. Probab.*, 8(3):539–575, 1980.
- [15] H. Tanaka. Lévy processes conditioned to stay positive and diffusions in random environments. volume 39 of *Adv. Stud. Pure Math.*, pages 355–376. Math. Soc. Japan, Tokyo, 2004.
- [16] K. Yano. Excursions away from a regular point for one-dimensional symmetric Lévy processes without Gaussian part. submitted, 2008.

- [17] K. Yano, Y. Yano, and M. Yor. On the laws of first hitting times of points for one-dimensional symmetric stable Lévy processes. submitted, 2008.
- [18] K. Yano, Y. Yano, and M. Yor. Penalising symmetric stable Lévy paths. *J. Math. Soc. Japan*, to appear.